

## WHIE TOUTHED STRAILES OF ANTERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

## pioneer Gi-Gred International, Inc.

Telliereas, there has been presented to the

#### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF eighteen years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic seed of the variety in a public repository as provided by LAW, the right to exclude others from selling the variety, or offering it for sale, or reproducing it, importing it, or exporting it, or using it in producing a hybrid or different ety therefrom, to the extent provided by the Plant Variety Protection Act at. 1542, as amended, 7 u.s.c. 2321 et seq.)

SOYBEAN

194111

In Eastimony Wincreot, I have hereunto set my hand and caused the seal of the Plant Variety Exotection Office to be affixed at the City of Washington, D. C. this 30th day of August in the year of our Lord one thousand nine hundred and ninety-one.

Col MAdig An Secretary of Agriculture

Allert

Konneth HEvan Commissioner

Plant Variety Protection Office Agricultural Marketing Service

U.S. DEPARTMENT				FORM APPROVED: OMB NO. 0681-0065				
AGRICULTURAL M.	ARKETING SERV	/ICE	A	pplication is required in a plant variety protec	n order to determine tion certificate is to			
APPLICATION FOR PLANT VARI	IETY PROTE	CTION CERTIFICA	TE b	e issued (7 U.S.C. 242 eld confidential until 7 U.S.C. 2426).	(1). Information is			
1. NAME OF APPLICANT(S)		2. TEMPORARY DESIGN	NATION 3	. VARIETY NAME	, , , , , , , , , , , , , , , , , , , ,			
Pioneer Hi-Bred International	, Inc.			9411				
4. ADDRESS (Street and No. or R.F.D. No., City, Sta	te, and Zip Code)	5. PHONE (Include area o		FOR OFFICIAL	USE ONLY			
700 Capital Square 400 Locust Street Des Moines, IA 50309		319-234-0335	ļ <sup>p</sup>	8900176				
6. GENUS AND SPECIES NAME	7. FAMILY NA	ME (Botanical)		DATE	11909			
Glycine Max	Legumin	osae		TIME TIME	A.M. P.M.			
8. KIND NAME	9.	DATE OF DETERMINAT	ION	AMOUNT FOR F	ILING			
Soybean		October, 1982	-	S J800				
30,504	January, 1987(incre							
10. IF THE APPLICANT NAMED IS NOT A "PERSO partnership, association, etc.)	N," GIVE FORM	OF ORGANIZATION (Co.		AMOUNTFORC	CERTIFICATE			
Corporation				\$ 200 mg 13	1991			
11. IF INCORPORATED, GIVE STATE OF INCORPORT	DRATION		1	2. DATE OF INCORP	ORATION			
13. NAME AND ADDRESS OF APPLICANT REPRES	SENTATIVE(S), I			TION AND RECEIVE	ALL PAPERS			
-Glark W. Jennings John Grace		Mary Helen M	15		61.			
3261 West Airline Highway 7301				- 400 Locust	Street			
Y <del>Water7oo, IA 50703-961</del> 0 P.O. 	1507KS 1570 N TA C	Des Moines,	TA DI nciude area d	0309 code):				
14. CHECK APPROPRIATE BOX FOR EACH ATTAC	CHMENT SUBMIT	TED						
a. X Exhibit A, Origin and Breeding History of	the Variety (See	Section 52 of the Plant Vo	riety Protec	ction Act.)				
b. 🔯 Exhibit B, Novelty Statement. c. 🖾 Exhibit C, Objective Description of Variet	ry (Peauest form	from Plant Variety Pratec	tion Office.	)	•			
d. Exhibit D, Additional Description of Variety		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		,				
e. X Exhibit E, Statement of the Basis of Appli	icant's Ownership							
15. DOES THE APPLICANT(S) SPECIFY THAT SEE SEED? (See Section 83(a) of the Plant Variety Pro		Yes (If "Yes,	" answer iter	ns 16 and 17 below)	X No			
16. DOES THE APPLICANT(S) SPECIFY THAT THIS LIMITED AS TO NUMBER OF GENERATIONS?	VARIETY BE	17. IF "YES" TO IT BEYOND BREE		ICH CLASSES OF PRO '	DDUCTION			
Yes X No		Foundation		Registered	Certified			
18. DID THE APPLICANT(S) PREVIOUSLY FILE	FOR PROTECT!	ON OF THE VARIETY IF	N THE U.S.7	Yes (i	If "Yes," give date)			
	•			ſŸ] No				
19. HAS THE VARIETY BEEN RELEASED, OFFER	RED FOR SALE,	OR MARKETED IN THE	U.S. OR O	THER COUNTRIES ?	If "Yes," give names			
		r ·		□ of co	untries and dates)			
20. The applicant(s) declare(s) that a viable samp			urnished w	لما	ind will be re-			
plenished upon request in accordance with su  The undersigned applicant(s) is (are) the own			alant variet	v. and believe(s) the	at the variety is			
distinct, uniform, and stable as required in Se Variety Protection Act.	ection 41, and is	entitled to protection u	nder the p	rovisions of Section	42 of the Plant			
Applicant(s) is (are) informed that false repre	esentation hereir	can jeopardize protecti	ion and res	<del></del>				
Clas b		april 6, 1989						
SIGNATURE OF APPLICANT	<del></del>			DATE	<del>/</del>			

FORM LS-470 (3-86) Attachment: 9411 Soybean (April, 1989)

Exhibit A:

Variety 9411 evolved from a cross of varieties (Williams 79 x A3127) X A3127. It is an F5-derived variety which was advanced to the F5 generation by modified single-seed descent. The F6 progeny row of 9411 was grown in Ohio during the summer of 1982. Subsequently, 9411 has undergone six years of extensive testing and purification and has been observed by the breeder to be uniform and stable for all plant traits from generation to generation, with no evidence of variants.

0.6 acres of 9411 (breeder's seed) were grown in 1987. 19 acres of parent seedstock (foundation seed equivalent) were grown in 1988.

Exhibit B: Variety 9411 is most similar to variety 9391. However, 9411 is significantly later maturing than 9391 (see Table 1.)

Exhibit E: Pioneer Hi-Bred International, Inc. is the sole, original, and first breeder of soybean variety 9411, for which it solicits a certificate of protection.

EXHIBIT C (Soybean)

U.S. DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE
LIVESTOCK, MEAT, GRAIN & SEED DIVISION
PLANT VARIETY PROTECTION OFFICE
BELTSVILLE, MARYLAND 20705

# OBJECTIVE DESCRIPTION OF VARIETY SOYBEAN (Glycine max L.)

	1	VARIETY NAME
NAME OF APPLICANT(S)	TEMPORARY DESIGNATION	
Pioneer Hi-Bred International, Inc.		9411
ADDRESS (Street and No., or R.F.D. No., City, State, and Zip Coo	le)	FOR OFFICIAL USE ONLY
Capital Square 400 Locust Street		PVPO NUMBER
Des Moines, IA 50309		8900176
Choose the appropriate response which characterizes the value in your answer is fewer than the number of boxes provided,	riety in the features described l place a zero in the first box w	below. When the number of significant digits hen number is 9 or less (e.g., 0 9).
1. SEED SHAPE:		
	, U	
2 L W	1   1	
1 = Spherical (L/W, L/T, and T/W ratios = $\langle 1.2 \rangle$ 3 = Elongate (L/T ratio $\rangle$ 1.2; T/W = $\langle 1.2 \rangle$		(L/W ratio > 1.2; L/T ratio = < 1.2) L/T ratio > 1.2; T/W > 1.2)
2. SEED COAT COLOR: (Mature Seed)		
1 = Yellow 2 = Green 3 = Brown	4 = Black 5 = Other	(Specify)
3. SEED COAT LUSTER: (Mature Hand Shelled Seed)		
2 1 = Dull ('Corsoy 79'; 'Braxton') 2 = Shiny ('Nebs	oy'; 'Gasoy 17')	
4. SEED SIZE: (Mature Seed)		
1 5 Grams per 100 seeds		
5. HILUM COLOR: (Mature Seed)		
6 1 = Buff 2 = Yellow 3 = Brown	4 = Gray 5 = Imperfect Bla	ck 6 = Black 7 = Other (Specify)
6. COTYLEDON COLOR: (Mature Seed)		
1 = Yellow 2 = Green		
7. SEED PROTEIN PEROXIDASE ACTIVITY:		3
2 1 = Low 2 = High		
8. SEED PROTEIN ELECTROPHORETIC BAND:		
1 = Type A (SP1 <sup>a</sup> ) 2 = Type B (SP1 <sup>b</sup> )		
9. HYPOCOTYL COLOR:		
1 = Green only ('Evans'; 'Davis') 2 = Green wit 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson';		Woodworth'; 'Tracy')
10. LEAFLET SHAPE:		
2 1 = Lanceolate 2 = Oval 3 = Ovate	4 = Other (Specify)	

11. LEA	FLET SIZE:	
2	1 = Small ('Amsoy 71'; 'A5312') 2 = Medium ('Corsoy 79'; 'Gasoy 17') 3 = Large ('Crawford'; 'Tracy')	
12. LEA	COLOR:	•
2	1 = Light Green ('Weber'; 'York') 2 = Medium Green ('Corsoy 79'; 'Braxton') 3 = Dark Green ('Gnome'; 'Tracy')	
13. FLO	VER COLOR:	
2	1 = White 2 = Purple 3 = White with purple throat	
14. POD	COLOR:	
1	1 = Tan 2 = Brown 3 = Black	
15. PLA	T PUBESCENCE COLOR:	
2	1 = Gray 2 = Brown (Tawny)	
16. PLAN	T TYPES:	
2	1 = Slender ('Essex'; 'Amsoy 71') 2 = Intermediate ('Amcor'; 'Braxton') 3 = Bushy ('Gnome'; 'Govan')	
17. PLAN	T HABIT:	
3	1 = Determinate ('Gnome'; 'Braxton') 2 = Semi-Determinate ('Will') 3 = Indeterminate ('Nebsoy'; 'Improved Pelican')	
L	3 = Indeterminate ('Nebsoy'; 'Improved Pelican')	
L	3 = Indeterminate ('Nebsoy'; 'Improved Pelican')  RITY GROUP:	
L	3 = Indeterminate ('Nebsoy'; 'Improved Pelican')	
18. MATU	3 = Indeterminate ('Nebsoy'; 'Improved Pelican')  RITY GROUP:  1 = 000	
18. MATU 0 7	3 = Indeterminate ('Nebsoy'; 'Improved Pelican')  RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC	3 = Indeterminate ('Nebsoy'; 'Improved Pelican')  RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC	3 = Indeterminate ('Nebsoy'; 'Improved Pelican')  RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC	RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC	3 = Indeterminate ('Nebsoy'; 'Improved Pelican')  RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC  0  0	RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC  0  0	RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC  0  0  FUNG	3 = Indeterminate ('Nebsoy'; 'Improved Pelican')  RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC  0  0  FUNG	RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC  0  0  FUNG	RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  0  0  0  FUNG  0	RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  BAC  0  0  FUNG  0	RITY GROUP:  1 = 000	
18. MATU  0 7  19. DISEA  0  0  0  0  0  0	RITY GROUP:  1 = 000	

19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) (Continued)												
	FUNGAL DISEASES: (Continued)											
	0	Pod and Stem Blight (Diaporthe phaseolorum var; sojae)										
	0	Purple Seed Stain (Cercospora kikuchii)		•								
	0 Rhizoctonia Root Rot (Rhizoctonia solani)											
		Phytophthora Rot (Phytophthora megasperma var. sojae)										
	2	Race 1 2 Race 2 2 Race 3 1	Race 4	2 Race 6 2 F	Race 7							
	2	Race 8 2 Race 9 2 Other (Specify)	10,11,13,15,17,2	1,23,24	· · · · · · · · · · · · · · · · · · ·							
	VIRAL DISEASES:											
	Bud Blight (Tobacco Ringspot Virus)											
	0	Yellow Mosaic (Bean Yellow Mosaic Virus)										
	0	Cowpea Mosaic (Cowpea Chlorotic Virus)										
	0	Pod Mottle (Bean Pod Mottle Virus)										
	0	Seed Mottle (Soybean Mosaic Virus)										
		ATODE DISEASES:										
		Soybean Cyst Nematode (Heterodera glycines)		·								
	1	Race 1 1 Race 2 1 Race 3 1	. Race 4 Other (	Specify)								
	Lance Nematode (Hoptolaimus Colombus)											
	$\vdash$	Northern Root Knot Nematode (Meloidogyne Hapla)										
	0	Peanut Root Knot Nematode (Meloidogyne arenaria)										
		Reniform Nematode (Rotylenchulus reniformis)	•		• .							
		OTHER DISEASE NOT ON FORM (Specify):										
0. P	HYSIOI	LOGICAL RESPONSES: (Enter 0 = Not Tested; 1 = Susce	ptible; 2 = Resistant)									
	0	Iron Chlorosis on Calcareous Soil										
		Other (Specify)										
. 16		REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = R	(ecietant)		· · · · · · · · · · · · · · · · · · ·							
	0	Mexican Bean Beetle (Epilachna varivestis)	esistant,									
		Potato Leaf Hopper (Empoasca fabae)	•									
		Other (Specify)	e e e e e e e e e e e e e e e e e e e									
		E WHICH VARIETY MOST CLOSELY RESEMBLES THA	AT SUBMITTED.		<del></del>							
	CHARA		CHARACTER	NAME OF VARIET	Y							
	ant Shap	0001	Seed Coat Luster	9441								
	ar Snapr		Seed Size	9391								
	af Size	9391	Seed Shape Seedling Pigmentation	9391								
	-	3331	occarry rightentation	9391								

## 23. GIVE DATA FOR SUBMITTED AND SIMILAR STANDARD VARIETY: Paired Comparison Data

VARIETY	NO. OF DAYS MATURITY	PLANT LODGING	CM PLANT HEIGHT	LEAFLET SIZE		SEED CON	ITENT	SEED SIZE G/100	NO. SEEDS/	
		SCORE		CM Width	CM Length	% Protein	% Oii	SEEDS	POD	
9411 Submitted	128.0	2.0	98.9			43.2	21.0	15.0		
9391 Name of Similar Variety	126.0	2.0	98.6	:		41.0	20.7	15.0		

### PUBLICATIONS USEFUL AS REFERENCE AIDS FOR COMPLETING THIS FORM:

- 1. Caldwell, B.E., ed. 1973. Soybeans: Improvement, Production, and Uses. Amer. Soc. Agron. Monograph No. 16.
- 2. Buttery, B.R. and R.I. Buzzell. 1968. Peroxidase activity in seeds of soybean varieties. Crop Sci., 8: 722-725.
- 3. Hymowitz, T. 1973. Electrophoretic analysis of SBTI-A2 in the USDA soybean germplasm collection. Crop Sci., 13: 420-421.
- 4. Payne, R.C. and L.F. Morris. 1976. Differentiation of soybean cultivars by seedling pigmentation patterns. J. Seed Technol. 1: 1-19.



## PIONEER HI-BRED INTERNATIONAL, INC. PLANT BREEDING DIVISION

DEPARTMENT OF SELF-POLLINATED CROPS
7301 NW 62ND AVENUE • P.O. BOX 85
JOHNSTON, 10WA 50131-0085
PHONE: (515) 270-3300
TELEFAX: (515) 253-2125

June 21, 1991

Jeffrey L. Strachan, Examiner Plant Variety Protection Office U.S. Department of Agriculture Room 500, NAL Building 10301 Baltimore Boulevard Beltsville, MD 20705

Subject: Soybean Application No. 8900176, '9411'

Dear Mr. Strachan:

Enclosed is an amended Exhibit B and relevant tables which provide evidence of novelty for variety '9411'. Comparisons are made against varieties '8628SE', '9331', '9391', 'CX326', 'CX366', and 'RA-501A'.

Variety '8628SE' has light tawny pubescence, whereas '9411' has the normal tawny pubescence. In addition, Dr. Garland (Research Director for Callahan Enterprises, Inc.) indicates the original PVP application shows '8628SE' has no specific resistance to Phytophthora. Variety '9411' has specific resistance to races 1-3 and 6-10 of Phytophthora.

Variety '9331' is an early to mid Group III, whereas '9411' is an early Group IV. Therefore, they are not tested together. However, data from 1989 was selected such that '9331', '9391' and '9411' were grown in the same locations, with '9331' and '9391' in one experiment at those locations and '9391' and '9411' in a different experiment. The data presented in Table 1 shows '9331' is 5 days earlier than '9391' at these selected locations. The probability of a greater t occurring by chance is less than 0.001. Data presented in Table 2 also shows '9391' is 2.3 days earlier than '9411'. The probability of a greater t occurring by chance is less than 0.05. Therefore, '9331' must be significantly earlier than '9441'.

Variety 'CX326' is 4.5 days earlier than '9411', based upon Table 3. The probability of a greater t occurring by chance is less than 0.001.

Variety 'CX366' and '9411' were not tested together. However, data from 1988 was selected such that 'CX366', '9391', and '9411' were grown in the same locations with 'CX366' and '9391' in one experiment at those locations, and '9391' and '9411' in a different experiment. Table 4 shows 'CX366' is 2.9 days earlier than '9391'. The probability of a greater t occurring by chance

Letter to Jeffrey L. Strachan June 21, 1991 Page 2

is less than 0.001. Data presented in Table 5 shows '9391' is 2.7 days earlier than '9411'. The probability of a greater toccurring by chance is less than 0.01. Therefore, 'CX366' must be significantly earlier than '9411'.

Variety 'RA-501A' is a Group V variety. As such, it is at least 10 days later than '9411'. Further, 'RA-501A' has specific resistance to Soybean Cyst Nematode race 3, whereas '9411' does not.

Finally, you had requested yearly breakouts of the comparison between '9391' and '9411'. While the data in Tables 2 and 5 (1989 and 1988 respectively) represents a subset of the information currently available, in the interest of brevity, I trust the significant differences found in these tables will suffice.

I hope the enclosed information is sufficient to allow you to issue a PVP Certificate on '9411'.

Sincerely,

John Grace

Assistant Project Manager

/sc Encl.

cc: Mary Helen Mitchell

Jim Miller

Table 1. Variety 9331 (X1) vs '9391' (X2) for maturity in days.

All observations are from plots planted using a randomized complete block design. Planted plot length was 21 feet, trimmed to 15 feet. Plot width was 4 30 inch rows, or 10 feet. Maturity was scored as the number of days from planting until 95% of the pods in the plot were mature. Data is presented for 1989. In our best scientific judgement this data results from plots treated in a statistically sound manner and as such, the use of the t test is completely valid.

REP	X1	x2	x1-x2	(x1-x2)2						
1	130	140	-10	100	SD**2=	0.75				
2 3	133	139	-6	36	SD=	0.86603				
. 3	117	121	-4	16	t=	-5.7735 ***				
4	118	120	-2	4	DF=	7				•
5	122	128		36						
4 5 6	121	126	-6 -5	.25	n=	8				
7 - 8	127	131	-4	16						
8	128	1.31	-3	9						
sum	996	1036	-40	242	ave đays	to maturity	for	9331	==	124.5
ave	124.5	129.5	5		ave days	to maturity	for	9391	=	129.5

<sup>\*\*\*</sup>t value required for significance at 0.1% level = 5.405, therefore probability of this t value occuring by chance < 0.001.

Table 2. Variety 9391 (X1) vs '9411' (X2) for maturity in days (1989).

All observations are from plots planted using a randomized complete block design. Planted plot length was 21 feet, trimmed to 15 feet. Plot width was 4 30 inch rows, or 10 feet. Maturity was scored as the number of days from planting until 95% of the pods in the plot were mature. Data is presented for 1989. In our best scientific judgement this data results from plots treated in a statistically sound manner and as such, the use of the t test is completely valid.

REP	X1	х2	x1-x2	(X1-X2)2						
1	137	139	-2	4	SD**2=	0.74777				
2	136	144	-8	64	SD=	0.86474				
2	121	122	-1	1	t=	-2.7465 *				
4	122	122	0	0	DF=	7				
5	123	126	-3	9	•					
4 5 6 7	124	126	-2	4	n=	8				
. 7	131	133	-2	4		•				
8	132	133	-1	1						
sum	1026	1045	-19	87	ave dave	to maturity	for	0301	_	128 3
ave	128.3 128.3			37		to maturity				

<sup>\*</sup>t value required for significance at 5% level = 2.365, therefore probability of this t value occuring by chance < 0.05.

Table 3. Variety CX326 (X1) vs '9411' (X2) for maturity in days.

All observations are from plots planted using a randomized complete block design. Planted plot length was 21 feet, trimmed to 15 feet. Plot width was 2 30 inch rows, or 5 feet. Maturity was scored as the number of days from planting until 95% of the pods in the plot were mature. Data is presented for 1988. In our best scientific judgement this data results from plots treated in a statistically sound manner and as such, the use of the t test is completely valid.

REP	x1	X2	X1-X2	(X1-X2)2									
1	127	132	-5	25	SD*	*2=		21667					
2	126	131	-5	25	SD=		0.4	46547					
- 3	127	133	-6	36	t=	-	-9	.6676	* * *				
4	128	135	-7	49	DF =			15	**				
4 5	128	136	8	64									
- 6	131	134	-3	9	n=			16					•
7	127	133	6	36									
8 9 10	129	135	-6	36									
9	125	129	-4	16									
10	124	126	-2	4		100		4					
11	125	128	3	9									
12	126	131	~5	25							•		
13	126	128	-2	4									
14	124	129	-5	25	*								
15	125	128	-3	9									
16	125	127	-2	4									
sum	2023	2095	-72	376	 ave	davs	t o	matuu	ritv	for	CX326	=	126.4
ave	126.4		-4.5		ave	days	to	matu	rity	for	9411	===	130.9

<sup>\*\*\*</sup>t value required for significance at 0.1% level = 4.073, therefore probability of this t value occuring by chance < 0.001.

Table 4. Variety CX 366 (X1) vs '9391' (X2) for maturity in days (1988).

All observations are from plots planted using a randomized complete block design. Planted plot length was 21 feet, trimmed to 15 feet. Plot width was 4 30 inch rows, or 10 feet. Maturity was scored as the number of days from planting until 95% of the pods in the plot were mature. Data is presented for 1988. In our best scientific judgement this data results from plots treated in a statistically sound manner and as such, the use of the t test is completely valid.

REP	<b>x</b> 1	<b>x</b> 2	x1-x2	(X1-X2)2					•		
1	127	130	-3	9	SD*	*2=	0.25444				
2	. 127	127	0	0	SD=		0.50442				
. 3	131	133	-2	4	t=		-5.7491	***			
4	120	126	-6	36	DF =		9				
5	122	126	-4	16							
4 5 6 7 8 9	137	141	-4	16	n=		10				
7	136	139	-3	9							
8	105	108	-3	9							
9	106	108	-2	4							
10	106	108	-2	.4							
sum	1217 121.7	1246 124.6	-29 -2.9	107			to matu			=	121.7 124.6

<sup>\*\*\*</sup>t value required for significance at 0.1% level = 4.781, therefore probability of this t value occuring by chance < 0.001.

Table 5. Variety 9391 (X1) vs '9411' (X2) for maturity in days (1988).

All observations are from plots planted using a randomized complete block design. Planted plot length was 21 feet, trimmed to 15 feet. Plot width was 4 30 inch rows, or 10 feet. Maturity was scored as the number of days from planting until 95% of the pods in the plot were mature. Data is presented for 1988. In our best scientific judgement this data results from plots treated in a statistically sound manner and as such, the use of the t test is completely valid.

REP	x1	X2	X1-X2	(X1-X2)2				
1	126	127	-1	1		SD**2=	0.62333	
2	129	129	0	0		SD=	0.78951	
3	134	139	-5	25		t=	-3.4198 **	
4	136	141	-5	25		DF =	9	
- 5	136	144	-8	64				
. 6	129	130	-1	1		n=	10	•
7	127	129	-2	4				
8 9	124	125	-1	1			•	
9	124	126	-2	4	*			
10	124	126	-2	. 4			·	
sum	1289	1316	-27	129		ave davs	to maturity	for 9391 = 128.9
ave	128.9	131.6	-2.7					for 9411 = 131.6

<sup>\*\*</sup>t value required for significance at 1% level = 3.250, therefore probability of this t value occurring by chance < 0.01.